

# **Application of vis-NIR spectroscopy for estimation of SOC and SOC fractions on soil samples burned under different laboratory conditions**

<sup>1</sup>Montorio, Raquel, <sup>2</sup>García Gil, Sandra, <sup>3</sup>Escuer Arregui, Marta, <sup>4</sup>Ortiz, Oriol, <sup>4</sup>Badía-Villas, David and <sup>1</sup>Pérez-Cabello, Fernando

<sup>1</sup>*Geoforest-IUCA research group. Department of Geography and Spatial Management. University of Zaragoza. Zaragoza, 50009, Spain*

<sup>2</sup>*Department of Geography and Spatial Management, University of Zaragoza, Zaragoza, 50009, Spain*

<sup>3</sup>*Technic School of Agricultural Engineering and Environmental Sciences, University of Zaragoza, Huesca, 22071, Spain*

<sup>4</sup>*Geoforest-IUCA research group. Technic School of Agricultural Engineering and Environmental Sciences, University of Zaragoza, Huesca, 22071, Spain*

## **Abstract**

Fire is one of the principal disturbances acting on forest soil organic carbon (SOC) (Nave et al., 2011). Fire affects both quantity and quality of SOC, with consequences depending on many fire characteristics such as duration or intensity. The impact on the SOC pools (i.e. labile, intermediate and recalcitrant pool) is also different, providing a complete vision of the real impact of fire on soils (Knicker, 2007). Laboratory analysis for the quantification of SOC and its fractions are complex and time-consuming. Alternatively, diffuse reflectance spectroscopy (DRS) has widely demonstrated its potential to estimate SOC with great accuracy in a time- and cost-effective way (Bellon-Maurel and McBratney, 2011). In this study we apply spectroscopic techniques in the vis-NIR spectral range to develop chemometric models (PLSR- partial least square regression) to estimate the oxidisable C and the three C pools. Surface (0-1 cm) and sub-surface (1-3 cm) samples (n=50) from soil monoliths burned under laboratory conditions at four levels of temperature and residence time, simulating different fire severity levels, were used. We also tested the performance of different spectral pre-treatments (i.e. derivative, absorbance transformation and normalization) for the development of the regression models. Results show the great capabilities of this technique, providing good estimates of oxidisable C and recalcitrant pool ( $R^2CV \sim 0.70$ ) and moderate results for the intermediate pool ( $R^2CV \sim 0.50$ ). The analysis of regression coefficients has also allowed the identification of spectral intervals for the prediction of SOC pools.

**Keywords:** soil carbon pools, spectroscopy, chemometric modeling

## References

Bellon-Maurel, V. McBratney, A. (2011). Near-infrared (NIR) and mid-infrared (MIR) spectroscopic techniques for assessing the amount of carbon stock in soils - Critical review and research perspectives. *Soil Biology and Biochemistry*, 43(7): 1398-1410. 10.1016/j.soilbio.2011.02.019

Knicker, H. (2007): How does fire affect the nature and stability of soil organic nitrogen and carbon? A review. *Biogeochemistry*, 85: 91-118. 10.1007/s10533-007-9104-4

Nave, L.E., Vance, E.D., Swanston, C.W., Curtis, P.S. (2011). Fire effects on temperate forest soil C and N storage. *Ecological Applications*, 21(4): 1189-1201. 10.1890/10-0660.1

**Acknowledgments:** This research was supported by the projects FUEGOPTIMO “Optimización del uso de quemas prescritas para la reducción del riesgo de incendios y la recuperación de pastos de alta biodiversidad en el Pirineo Central” (UZ2020-TEC-01) and PaF “Análisis dinámico de la resiliencia de los Paisajes Forestales afectados por el Fuego mediante indicadores espectrales multisensor” (PID2020-118886RB-I00, Ministerio de Ciencia e Innovación).