

Future Climate Change Impact on Wildfire Danger over the Mediterranean: the case of Greece

¹Rovithakis Anastasios, ¹Voulgarakis Apostolos, ¹Grillakis Manolis, ¹Seiradakis Kostadinos, ²Giannakopoulos Christos, ²Karali Anna, ³Field Robert and ⁴Lazaridis Mihalís

¹*School of Environmental Engineering, Technical University of Crete, Chania, Greece and Leverhulme Centre for Wildfires, Environment and Society, Imperial College London, London, UK*

²*Institute for Environmental Research and Sustainable Development, National Observatory of Athens, Athens, Greece*

³*Department of Applied Physics and Applied Mathematics, Columbia University, New York, USA*

⁴*School of Environmental Engineering, Technical University of Crete, Chania, Greece*

Abstract

Recent studies have shown that temperature and precipitation in the Mediterranean are expected to change, indicating longer and more intense summer droughts that even extend out of season. In connection to this, the frequency of forest fire occurrence and intensity will likely increase. In the present study, the changes in future fire danger conditions are assessed for the different regions of Greece using the Canadian Fire Weather Index (FWI). Gridded future climate output as estimated from six regional climate models from the Coordinated Regional Downscaling Experiment (CORDEX) are utilized. We use three Representative Concentration Pathways (RCPs), consisting of an optimistic emissions scenario where emissions peak and decline beyond 2020 (RCP2.6), a mid-of-the-road scenario (RCP4.5) and a pessimistic scenario, in terms of mitigation, where emissions continue to rise throughout the century (RCP8.5). The FWI projections were assessed for two future time periods, 2021-2050 and 2071-2100, comparing to a reference time period in the recent past 1971-2000. Based on established critical fire risk threshold values for Greece, the future change in days with critical fire risk were calculated for different Greek domains. The results show that future fire danger is expected to progressively increase in the future, especially in the high-end climate change scenario, with southern and eastern regions of Greece exhibiting increases in the FWI that exceed 20 FWI units, on average. Furthermore, southern Crete, the Aegean Islands, the Attica region, as well as parts of eastern and southern Peloponnese are predicted to experience a larger increase in the fire danger, with an additional 12-17 potential fire days in the distant future (2070-2100) when compared to the reference period, under the RCP8.5 scenario.

Keywords: Fire danger, fire weather index, Euro-CORDEX, wildfires

Acknowledgments: This project is supported by the project/program \"National Network on Climate Change and its Impacts - Climact\" financed by the Public Investment Program of Greece and based on the results from FWI simulations provided by Copernicus Climate Change Service. Also, this research was partially funded by the Leverhulme Centre for Wildfires, Environment, and Society through the Leverhulme Trust

DRAFT