

# **A reconstruction of fire and vegetation in Cantabria, northern Iberian Peninsula throughout the Holocene using a multi-proxy approach**

<sup>1</sup>Braunthal, Ashley Nicole, <sup>1</sup>Pèlachs Mañosa, Albert, <sup>1</sup>Cunill Artigas, Raquel, <sup>2</sup>Sanchez Morales, Marc and <sup>3</sup>Perez Obiol, Ramon

<sup>1</sup>*Department of Geography, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, 08193, Spain*

<sup>2</sup>*Department of Biosciences, Universitat de Vic-Universitat Central de Catalunya, Vic, Barcelona, 08500, Spain*

<sup>3</sup>*Department of Animal Biology, Plant Biology and Ecology, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, 08193, Spain*

## **Abstract**

Sedimentary palaeofire records are critical to our understanding of past ecology, fire, climate, and human-land relationships in postglacial quaternary environments. This data provides insights into the favourable environmental (e.g., flora and vegetation) and climate (e.g., hydrological) conditions that impact fire propagation, intensity, and frequency.

Here we present a study of a multi-proxy approach to reconstruct fires in Cantabria throughout the Holocene from a peat record site (Riófrío peatbog, Vega de Liébana, Cantabria in the northern Iberian Peninsula, 43.037°N, 4.697 W°, 1740 MASL). Besides contributing valuable palaeoecological knowledge, our work will fill in research gaps of postglacial quaternary environmental history of the Iberian peninsula's mountain regions (Ruiz-Fernandez et al., 2016) and validate charcoal classification methods.

Macroscopic and microscopic charcoal concentrations in peat records are established palaeofire proxies. During a fire, charcoal particles undergo primary deposition via aeolian fallout, with the size decreasing over distance (Whitlock & Larsen, 2005). The concentration and deposition rate of particulate charcoal, through counting and size classification, can signal the timing, intensity, patterns, and proximity of the fires to the deposition site. The morphology (i.e., qualitative particle identity classification) and morphometry of the particles (i.e., length:width ratio calculations) provide information on the kind of flora that burned (Enache and Cumming 2006; Mustaphi and Pisaric, 2014; Umbanhowar and McGrath, 1998).

As such, we have combined these approaches: at a 1 cm resolution, we applied the 27

classifications method from Mustaphi and Pisaric (2014) and placed them into size classes; we also determined the length:width ratios and organic matter content (loss on ignition, LOI). Additionally, we looked at pollen and non-pollen palynomorphs (NPP) to further interpret and validate signals of past climate, dominant burned flora, and surrounding vegetation.

From this multi-proxy data, aside from a fire reconstruction, we will distinguish periods of open versus forested vegetation (i.e. maintenance fires versus newly burned from charcoal and pollen data; Fyfe et al., 2003; Pérez-Obiol et al., 2012; Rodríguez-González et al., 2023). The study allows us to validate secondary charcoal deposition through erosion proxies (i.e., NPP). Organic matter data (i.e., LOI) could provide us with indirect connections between fire dynamics and large-scale climate processes (i.e., flooding, Ishii et al., 2017).

**Keywords:** fire, palaeoclimate, particulate charcoal, landscape

### References

1. Enache, M.D., and Cumming, B.F., 2009, Extreme fires under warmer and drier conditions inferred from sedimentary charcoal morphotypes from Opatcho Lake, central British Columbia, Canada: *The holocene*, v. 19, p. 835-846, doi:10.1177/0959683609337357.
2. Fyfe, R.M., Brown, A.G., and Rippon, S.J., 2003, Mid- to late-Holocene vegetation history of Greater Exmoor, UK: estimating the spatial extent of human-induced vegetation change: *Vegetation history and archaeobotany*, v. 12, p. 215-232, doi:10.1007/s00334-003-0018-3.
3. Ishii, Y., Hori, K., and Momohara, A., 2017, Middle to late Holocene flood activity estimated from loss on ignition of peat in the Ishikari lowland, northern Japan: *Global and planetary change*, v. 153, p. 1-15, doi:10.1016/j.gloplacha.2017.04.004.
4. Mustaphi, C.J.C., and Pisaric, M.F.J., 2014, A classification for macroscopic charcoal morphologies found in Holocene lacustrine sediments: *Progress in physical geography*, v. 38, p. 734-754, doi:10.1177/0309133314548886.
5. Pérez-Obiol, R., Bal, M.-C., Pèlachs, A., Cunill, R., and Soriano, J.M., 2012, Vegetation dynamics and anthropogenically forced changes in the Estanilles peat bog (southern Pyrenees) during the last seven millennia: *Vegetation history and archaeobotany*, v. 21, p. 385-396, doi:10.1007/s00334-012-0351-5.
6. Rodríguez-González, J.-M., Sánchez-Morales, M., Nadal-Tersa, J., Pèlachs, A., and Pérez-Obiol, R., 2023, Paleoenvironmental reconstruction for the last 3500 years in the southern Pyrenees from a peat bog core in Clots de Rialba: *Diversity*, v. 15, p. 390, doi:10.3390/d15030390.
7. Ruiz-Fernández, J., Oliva, M., Cruces, A., Lopes, V., Freitas, M. da C., Andrade, C., García-Hernández, C., López-Sáez, J.A., and Gerales, M., 2016, Environmental evolution in the Picos de Europa (Cantabrian Mountains, SW Europe) since the Last Glaciation: *Quaternary science reviews*, v. 138, p. 87-104, doi:10.1016/j.quascirev.2016.03.002.
8. Umbanhowar, C.E., Jr, and Mcgrath, M.J., 1998, Experimental production and analysis of microscopic charcoal from wood, leaves and grasses: *The holocene*, v. 8, p. 341-346, doi:10.1191/095968398666496051.
9. Whitlock, C., and Larsen, C., 2005, Charcoal as a fire proxy, in *Tracking Environmental Change Using Lake Sediments*, Dordrecht, Kluwer Academic Publishers, p. 75-97.