

Plant species impact on runoff and soil erosion in a Mediterranean shrubland

¹Cerda, A, ²Javier Cerdà Benito and ³Francisco Escriva-Saneugenio

¹*Soil Erosion and Degradation Research Group. Department of Geography, Valencia University, Blasco Ibàñez, 28, 46010 Valencia, Spain.*

²*Escuela Técnica Superior de Ingeniería Agronómica y del Medio Natural. Univeristat Politècnica de València. Camino de Vera s/n. 46022 Valencia, Spain.*

³*Soil Erosion and Degradation Research Group, <https://soilerosion.eu>. Department of Geography, Valencia University, Blasco Ibàñez, 28, 46010 Valencia, Spain.*

Abstract

Shrubland is a Mediterranean biome characterized by densely growing evergreen shrubs adapted to fire events. To date, scientific research has focused on the impact of vegetation on soil erosion mainly through the control that plant biomass or plant cover exerts on sediment delivery and runoff discharge, being the individual plant species' influence on hydrological and erosional processes not achieved in detail. The objective of this research is to determine: i) runoff and soil losses in shrubland-covered rangeland at Sierra de Enguera, Spain; and ii) how four plant species affect soil and water losses. We measured soil cover, soil properties, runoff discharge and sediment yield under natural rainfall for five years (2010-2014) in a typical shrubland burnt in 1999. Four plant species were selected with 4 plots each: *Ulex parviflorus* Pourr., *Pistacia lentiscus* L., *Quercus coccifera* L., and *Rosmarinus officinalis* L. Despite that the soil properties and plant cover did not exhibit statistically significant differences among plant species, the runoff discharge was lower on *Q. coccifera* (4.87 %, SE 0.24) and *P. lentiscus* (6.24 %, SE 0.51) than on *U. parviflorus* (13.41 %, SE 0.58) and *R. officinalis* (13.84 %, SE 1.23). Sediment concentrations were, respectively, 3.91, 4.33, 4.31, and 4.88 g l⁻¹, and the differences between *R. officinalis* and the other species were statistically significant. The runoff discharge determined differences in soil erosion rates amongst the plant species with lower rates on *P. lentiscus* (1.36 Mg ha⁻¹ y⁻¹) and *Q. coccifera* (1.53 Mg ha⁻¹ y⁻¹), than on *U. parviflorus* (3.17 Mg ha⁻¹ y⁻¹) and *R. officinalis* (3.85 Mg ha⁻¹ y⁻¹). This long-term in situ study indicated that *Q. coccifera* and *P. lentiscus* are more efficient in controlling runoff discharge and soil losses than *U. parviflorus* and *R. officinalis* one decade after a fire. We discuss these results in light of the recent findings by the scientific community on the role of the canopy cover (rainfall interception), soil macropore and root system, and the water repellency that controls the hydrological response of the soil (e.g. runoff generation, infiltration). The information supplied by 5 years of research is relevant for restoration and rehabilitation programs and advises that *Q. coccifera* and *P. lentiscus* are

the most efficient plant species to control soil and water losses within the Mediterranean shrubland. This is an applied science approach for the better management of rangelands.

Keywords: Plants, Shrubland, Plots, Rainfall, Runoff, Soil erosion, Sediment; Mediterranean

References

Cerdà, A., Lucas-Borja, M. E., Franch-Pardo, I., Úbeda, X., Novara, A., López-Vicente, M., ... & Pulido, M. (2021). The role of plant species on runoff and soil erosion in a Mediterranean shrubland. *Science of The Total Environment*, 799, 149218. Rodrigo Comino, J., Keesstra, S. D., & Cerdà, A. (2018). Connectivity assessment in Mediterranean vineyards using improved stock unearthing method, LiDAR and soil erosion field surveys. *Earth Surface Processes and Landforms*, 43(10), 2193-2206. Keesstra, S., Nunes, J. P., Saco, P., Parsons, T., Poepl, R., Masselink, R., & Cerdà, A. (2018). The way forward: can connectivity be useful to design better measuring and modelling schemes for water and sediment dynamics?. *Science of the Total Environment*, 644, 1557-1572. Cerdà, A., Borja, M. E. L., Úbeda, X., Martínez-Murillo, J. F., & Keesstra, S. (2017). *Pinus halepensis* M. versus *Quercus ilex* subsp. *Rotundifolia* L. runoff and soil erosion at pedon scale under natural rainfall in Eastern Spain three decades after a forest fire. *Forest ecology and management*, 400, 447-456. Cerdà, A., Novara, A., Dlapa, P., López-Vicente, M., Úbeda, X., Popović, Z., Mekonnen, M., Terol, E., Janizadeh, S., Mbarki, S., Saldanha Vogelmann, E., Hazrati, S., Sannigrahi, S., Parhizkar, M., Giménez-Morera, A. 2021. Rainfall and water yield in Macizo del Caroig, Eastern Iberian Peninsula. Event runoff at plot scale during a rare flash flood at the Barranco de Benacancel. *Cuadernos de Investigación Geográfica* 47, <http://doi.org/10.18172/cig.4833>.

Acknowledgments: This research was funded by REACT4MED: Inclusive Outscaling of Agro-Ecosystem Restoration Actions for the Mediterranean. REACT4MED Project (Grant Agreement No. 2122) funded by PRIMA, Horizon 2020, European Union's Framework Programme for Research and Innovation info@react4med.eu. SECOMAL AICO/2021/68 Soil Erosion CONTROL in Mediterranean Agriculture Land. Conselleria d'Innovació, Universitats, ciència i societat digital. Firelinks EU Cost Action 18135.